

Name: _____ D.O.B.: _____

Allergic to: _____

 Weight: _____ lbs. Asthma: ☐ **Yes (higher risk for a severe reaction)** ☐ **No**

**PLACE
PICTURE
HERE**

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.

Extremely reactive to the following allergens: _____

THEREFORE:

- ☐ If checked, give epinephrine immediately if the allergen was **LIKELY** eaten, for **ANY** symptoms.
- ☐ If checked, give epinephrine immediately if the allergen was **DEFINITELY** eaten, even if no symptoms are apparent.

FOR **ANY** OF THE FOLLOWING:
SEVERE SYMPTOMS



LUNG

Shortness of breath, wheezing, repetitive cough



HEART

Pale or bluish skin, faintness, weak pulse, dizziness



THROAT

Tight or hoarse throat, trouble breathing or swallowing



MOUTH

Significant swelling of the tongue or lips



SKIN

Many hives over body, widespread redness



GUT

Repetitive vomiting, severe diarrhea



OTHER

Feeling something bad is about to happen, anxiety, confusion

**OR A
COMBINATION**
of symptoms
from different
body areas.

1. **ADMINISTER EPINEPHRINE IMMEDIATELY.**
2. **Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
 - Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
 - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
 - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
 - Alert emergency contacts.
 - Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS



NOSE

Itchy or runny nose, sneezing



MOUTH

Itchy mouth



SKIN

A few hives, mild itch



GUT

Mild nausea or discomfort

**FOR MILD SYMPTOMS FROM MORE THAN ONE
SYSTEM AREA, GIVE EPINEPHRINE.**

**FOR MILD SYMPTOMS FROM A SINGLE SYSTEM
AREA, FOLLOW THE DIRECTIONS BELOW:**

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

Epinephrine Dose: ☐ 0.1 mg IM (intramuscular) ☐ 0.15 mg IM
☐ 0.3 mg IM ☐ 1 mg IN (intranasal) ☐ 2 mg IN

Antihistamine Brand or Generic: _____


Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____

☐ Patient may self-carry ☐ Patient may self-administer


HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

1. Remove Auvi-Q from the outer case. Pull off red safety guard.
2. Place black end of Auvi-Q against the middle of the outer thigh.
3. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
4. Call 911 and get emergency medical help right away.




HOW TO USE EPIPEN®, EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, VIATRIS AUTO-INJECTOR, VIATRIS

1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, remove the blue safety release by pulling straight up.
3. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
4. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.




HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENALCLICK®), USP AUTO-INJECTOR, AMNEAL PHARMACEUTICALS

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip. Grasp the auto-injector in your fist with the red tip pointing downward.
3. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh. Press down hard and hold firmly against the thigh for approximately 10 seconds.
4. Remove and massage the area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES

1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, pull off the blue safety release.
3. Place the orange tip against the middle of the outer thigh at a right angle to the thigh.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

HOW TO USE NEFFY® (EPINEPHRINE NASAL SPRAY)

1. Remove neffy from packaging. Pull open the packaging to remove the neffy nasal spray device.
2. Hold device as shown. Hold the device with your thumb on the bottom of the plunger and a finger on either side of the nozzle. Do not pull or push on the plunger. Do not test or prime (pre-spray). Each device has only 1 spray.
3. Insert the nozzle into a nostril until your fingers touch your nose. Keep the nozzle straight into the nose pointed toward your forehead. Do not point (angle) the nozzle to the nasal septum (wall between your 2 nostrils) or outer wall of the nose.
4. Press plunger up firmly until it snaps up and sprays liquid into the nostril. Do not sniff during or after the dose is given. If any liquid drips out of the nose, you may need to give a second dose of neffy after checking for symptoms.
5. If symptoms don't improve or worsen within 5 minutes of initial dose, administer a second dose into the same nostril with a new neffy device.



Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can worsen quickly.

EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: _____

DOCTOR: _____ PHONE: _____

PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____ PHONE: _____

NAME/RELATIONSHIP: _____ PHONE: _____

NAME/RELATIONSHIP: _____ PHONE: _____

FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN & TREATMENT AUTHORIZATION

Appendix F-4

EPINEPHRINE AUTHORIZATION & ANTIHISTAMINE AUTHORIZATION FOR USE WITH ALLERGY ACTION PLAN Release and indemnification agreement

PART I TO BE COMPLETED BY PARENT OR GUARDIAN

☐ I hereby request designated school personnel to administer **epinephrine** as directed by this authorization. I agree to release, indemnify, and hold harmless the designated school personnel, or agents from lawsuits, claim expense, demand or action, etc., against them for administering this medication, provided the designated school personnel comply with the Licensed Healthcare Provider (LHCP) or parent or guardian orders set forth in accordance with the provision of the attached Food Allergy and Anaphylaxis Care Plan. I am aware that the medication may be administered by a specifically trained non-health professional. I have read the procedures outlined on the back of this form and assume responsibility as required. I understand that emergency medical services (EMS) will always be called when epinephrine is given, whether or not the student manifests any symptoms of anaphylaxis. Two pre-measured doses will be needed in school.

☐ I hereby request designated school personnel to administer **antihistamine and/or inhaled medication** as directed by this authorization. I agree to release, indemnify, and hold harmless the designated school personnel, or agents from lawsuits, claim expense, demand or action, etc., against them for helping this student use medication, provided the designated school personnel comply with the Licensed Healthcare Provider (LHCP) or parent or guardian orders set forth in accordance with the provision of the **attached** Food Allergy and Anaphylaxis Care Plan. I have read the procedures outlined below this form and assume responsibility as required.

Student Name (Last, First, Middle)

Date of Birth

Allergies:

School:

School Year:

PART II SEE PAGE 1 OF FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN & TREATMENT AUTHORIZATION – Completed by Parent/Guardian and Student, if applicable

The injectable epinephrine dosage will be given as noted and detailed on the attached Allergy Action Plan

Check ☒ the appropriate boxes:

- ☐ Allergy Action Plan is attached with orders signed by Licensed Healthcare Provider
- ☐ It is not necessary for the student to carry his/her medication during school, the prescribed medication will be kept in the clinic or other approved school location.
- ☐ The student is to carry epinephrine during school and school sanctioned events with principal/school nurse approval. (An additional epinephrine, to be used as backup, is advised to be kept in the clinic or other approved school location and Appendix F-21B is signed) Additionally, I believe that this student has received information on how and when to use the epinephrine and that he or she demonstrates its proper use.
- ☐ The antihistamine medication will be given as noted and detailed on the attached Allergy Action Plan, if applicable.
- ☐ The inhaled medication will be given as noted and detailed on the attached Allergy Action Plan, if applicable.

Parent or Guardian Name (Print or Type)

Parent or Guardian (Signature)

Telephone

Date

Student Name (Print or Type)

Student Signature (Required if Self Carry in addition to Appendix F-21B)

Date

PART III TO BE COMPLETED BY LICENSED NURSE OR TRAINED ADMINISTRATOR OF MEDICATION

Check ☒ as appropriate:

- ☐ Part I and II are completed and signed.
- ☐ Food Allergy and Anaphylaxis Care Plan is completed in its entirety and signed by the LHCP and attached to this form.
- ☐ Epinephrine, Antihistamine and Inhaled Medication, if applicable, are appropriately labeled.
- ☐ I have reviewed the proper use of their epinephrine with the student and, ☐ agree ☐ disagree that student should self carry their device in school. Appendix F-21B is also reviewed and attached.
- ☐ If self-carry and parent does not supply 2nd dose of epinephrine for clinic, parent must sign acknowledge and refusal to send medication form, Appendix F-25.

_____ Date any unused medication was collected by the parent or properly disposed. (Within one week after expiration of the physician order or on the last day of school).

Signature

Date

PARENT INFORMATION ABOUT MEDICATION PROCEDURES

1. **In no case may any health, school, or staff member administer any medication outside the framework of the procedures outlined here in the *Office of Catholic Schools Policies and Guidelines* and *Virginia School Health Guidelines* manual.**
2. **Schools do NOT provide routine medications for student use.**
3. Medications should be taken at home whenever possible. The first dose of any new medication must be given at home to ensure the student does not have a negative reaction.
4. Medication forms are required for each Prescription and Over the Counter (OTC) medication administered in school.
5. **All medication taken in school must have a parent/guardian signed authorization. Prescription medications, herbals and OTC medications taken for 4 or more consecutive days also require a licensed healthcare provider's (LHCP) written order. No medication will be accepted by school personnel without the accompanying complete and appropriate medication authorization form.**
6. **The parent or guardian must transport medications to and from school.**
7. Medication must be kept in the school health office, or other principal approved location, during the school day. All medication will be stored in a locked cabinet or refrigerator, within a locked area, accessible only to authorized personnel, unless the student has prior written approval to self-carry a medication (e.g. inhaler, auto-injector). If the student self carries, it is advised that a backup medication be kept in the clinic. If a backup auto-injector is not supplied, please complete Appendix F-25.
8. Parents/guardians are responsible for submitting a new medication authorization form to the school at the start of the school year and each time there is a change in the dosage or the time of medication administration.
9. A Licensed Health Care Provider (LHCP) may use office stationery, prescription pad or other appropriate documentation in lieu of completing the Food and Anaphylaxis Care Plan. The following information written in lay language with no abbreviations must be included and attached to this medication administration form. Signed faxes are acceptable.
 - a. Student name
 - b. Date of Birth
 - c. Diagnosis
 - d. Signs or symptoms
 - e. Name of medication to be given in school
 - f. Exact dosage to be taken in school
 - g. Route of medication
 - h. Time and frequency to give medications, as well as exact time interval for additional dosages.
 - i. Sequence in which two or more medications are to be administered
 - j. Common side effects
 - k. Duration of medication order or effective start and end dates
 - l. LHCP's name, signature and telephone number
 - m. Date of order
10. All prescription medications, including physician's samples, must be in their original containers and labeled by a LHCP or pharmacist. Medication must not exceed its expiration date.
11. All Over the Counter (OTC) medication must be in the original, small, sealed container with the name of the medication and expiration date clearly visible. Parents/guardians must label the original container of the OTC with:
 - a. Name of student
 - b. Exact dosage to be taken in school
 - c. Frequency or time interval dosage is to be administered
12. The student is to come to the clinic or a predetermined location at the prescribed time to receive medication. Parents must develop a plan with the student to ensure compliance. Medication will be given no more than one half hour before or after the prescribed time.
13. **Students are NOT permitted to self medicate. The school does not assume responsibility for medication taken independently by the student.** Exceptions may be made on a case-by-case basis for students who demonstrate the capability to self-administer emergency life saving medications (e.g. inhaler, auto injector)
14. Within one week after expiration of the effective date on the order, or on the last day of school, the parent or guardian must personally collect any unused portion of the medication. Medications not claimed within that period will be destroyed.