Emergency Care Plan



Student:	Grade:	School Contac	t:	DOB:
Asthmatic: Yes	No (increased risk for sev	rere reaction Allergen(s	s):	
Mother:		_MHome #:	_MWork #:	MCell #:
Father:		_ FHome #:	_ FWork #:	FCell #:
Emergency Contact:		Relationship:	Phone:	
 MOUTH THROAT SKIN STOMACH LUNG HEART Th 	Itching, tightness in thro	s, tongue or mouth, mout at, hoarseness, cough ag of face and extremities aps, vomiting, diarrhea etitive cough, wheezing ag out" s can change quickly	h "feels hot"	Student Photo
STAFF MEMBERS IN	STRUCTED: Administration	☐ Classroom Teacher(s)☐ Support Staff	☐ Special Area T ☐ Transportation	. ,
TREATMENT:	Rinse contact area with	water if appropriate		
Treatment should be initiated with symptoms without waiting for symptoms Benadryl ordered: Yes No				
IF INGESTION OR SUSPECTED INGESTION OF ALLERGEN OCCURS, SYMPTOMS ARE PRESENT AND EPINEPHRINE IS ORDERED, GIVE EPINEPHRINE IMMEDIATELY AND CALL 911. Preferred Hospital if transported: Epinephrine provides a 20 minute response window. After epinephrine, a student may feel dizzy or have an increased heart rate. This is a normal response. Students receiving epinephrine should be transported to the hospital by ambulance. A staff member should accompany the student to the emergency room if the parent, guardian or emergency contact is not present and adequate supervision for other students is present.				
	☐ Medication available or			
Healthcare Provider: Si	gnature		Phone:	
Written by:			Date:	
	☐ Copy provided to Pare		sent to Healthcare Provide	
Parent/Guardian Sign	nature to share this plan w	ith Provider and School S	taff:	