Individual Emergency Health Plan for Anaphylaxis

PICTURE						
OF						
STUDENT						

STUDENT STUDENT								
Name:	Allergic to:							
D/O/B:								
Weight:								
	Class:				. 1. 5- 3			
Asthmatic (Check box if YES) Student has an increased risk of a severe allergic reaction. Epinephrine should be given <u>first</u> (before asthma medications) in case of a reaction with any breathing symptoms.								
« STEP 1 TREATMENT »								
SIG	INS OF AN ALLERGIC REACTION	MEDICATION						
		(indicate medication name/dose/route, to be determined by physician authorizing treatment)						
Category	Symptom(s)	Epinep	hrine	Antihista	amine			
	No symptoms and suspected ingestion of	First Dose		Yes 🗆				
	allergen.	Second Dose		No 🗆				
	No symptoms and known ingestion of allergen.	First Dose		Yes				
		Second Dose		No 🗆				
Mouth	Itching, tingling, or swelling of lips, tongue, or	First Dose		Yes □				
	mouth	Second Dose		No 🗆				
Nose/Eyes	Hay fever-like symptoms: runny, itchy nose; red eyes	First Dose		Yes \square				
		Second Dose		No 🗆				
Skin(1)	Localized hives and/or localized itchy rash	First Dose		Yes				
		Second Dose		No 🗆	Establish to			
Skin(2)	Hives and/or itchy rash on more than one part	First Dose		Yes				
	of the body, swelling of face or extremities			No 🗆				
Gut	Nausea, abdominal cramps, vomiting, diarrhea	First Dose		Yes 🗆				
		Second Dose		No 🗆				
Throat	pat Hacking cough, tightening of throat,			Yes 🗆	*			
	hoarseness, difficulty swallowing	Second Dose		No 🗆				
Lung	Shortness of breath; wheezing; short, frequent, shallow cough	First Dose		Yes 🗆				
		Second Dose		No 🗆				
Heart	Weak pulse, low blood pressure, fainting, dizzy, pale, cyanosis (blueness)	First Dose		Yes □				
		Second Dose		No 🗆				
Multiple	Symptoms from two or more of the above	First Dose		Yes 🗆				
	categories.	Second Dose		No 🗆				
DOSAGE								
Epinephrine: Brand Name: Dosage: () 0.15 mg IM () 0.3 mg IM								
Antihistami	ne Medication: Brand Name:		Dosage:					
Inhaler-Bronchodilator Brand Name: Dosage:								
Student may self-carry epinephrine: Yes No Student may self-administer epinephrine: Yes No								

« After administering treatment, turn page over for EMERGENCY CONTACTS »

«2. EMERGENCY CONTACTS»

				**					
(4.1	NAME	RELATIONSHIP	PHONE	INSTRUCTIONS					
			NUMBER	- 911 is the <u>first</u> call that must be made					
1									
	911			after administering epinephrine.					
				- Indicate to the first responders that the					
				student is suffering from an allergic					
				reaction and may require additional					
				epinephrine.					
2	Physician:								
		Student's allergist or							
	Dr	pediatrician							
3	Parent/Guardian:	(Specify Relationship):	100						
			K.						
4	Parent/Guardian:	(Specify Relationship):							
-									
	E Contact	(Specify Relationship):							
5	Emergency Contact	(Specify Relationship).							
	(name):								
6	If Possible - What hospital								
0	would you like the child		10-12-5						
-	transported to in case of								
	an allergic reaction?		1 - 1 - 1 - 1						
Г									
		Administration	of Epinephi	rine					
<u></u>									
D	ate:		Who ada	ninistered the epinephrine?					
D	osage:								
Time:									
T	he forenamed student is my p	patient and I have authori	zed the treatm	nent protocol outlined on the preceding					
	age and affirm that there are								
Physician signature and date:									
I	I authorize the administration of epinephrine, antihistamine or other specified medication to the forenamed								
student as ner the treatment protocol outlined on the preceding page									

Parent/Guardian signature and date: